The Bryologist 95(4), 1992, pp. 492–501
Copyright © 1992 by the American Bryological and Lichenological Society, Inc.

Index to Volume 95 of THE BRYOLOGIST[1]

CYNTHIA M. GALLOWAY
Department of Biology, Texas A&I University, Kingsville, TX 78363

Abietinella abietina, 31
Abrothallus parmeliarum, 404; welwitzschii, 33+
Acarospora badiofusca, 33; fuscata, 139; glaucocarpa, 393; pelicypha, 393; schleicheri, 330; sinopica, 33, 393
Acrobolbus ciliatus, 309; rhizophyllus, 310
Adelanthus decipiens, 289
Adelolecia pilati, 393
Adelotheciaceae, 287
Adelothecium bogotense, 287+*
Adenostoma, 330; fasciculatum, 330
Aerobryidium subpiligerum, 289
Agrestia hispida, 33
Agyrophora lyngei, 393
Alaska, Calycularia crispula new to, 266; lichens of the Tuxedni Wilderness Area, 20
Alectoria nigricans, 24, 393; ochroleuca, 24, 393; vexillifera, 394
alectoronic acid, 2, 378+
alkaline phosphatase, 321+
Allantoparmelia almquistii, 394; alpicola, 394
ALLEN, B. [Review:] H. Crum. Liverworts and Hornworts of Southern Michigan, 374
ALLEN, B. [Review:] J.-P. Frahm. Campylopodes Centrali-Africanae, 371
ALLEN, B. [Review:] J.-P. Frahm. Campylopodes Exsiccatae Fasc. VII, Nr. 151–175, 371
ALLEN, B. [Review:] S. R. Gradstein. Bryophyta Neotropica Exsiccata. Fascicle VI, 371
ALLEN, B. *See* MAGILL, R. E. & B. ALLEN, 128, 363
ALLEN, B. Teratological axillary "hairs" in Didymodon tophaceus, 97
ALLEN, B. & R. E. MAGILL. Recent literature on mosses—152, 250; —154, 478
Allisoniaceae, 268
Aloina bifrons, 280
Alophosia, 272
Alsia californica, 69+
ALSTRUP, V. *See* ISBRAND, S. & V. ALSTRUP, 233
Amandinea, 436
Amblyolejeunea, 42
Amblystegium, 334
Ames, M. P., 73
Amygdalaria consentiens, 394; elegantior, 392+; panaeola, 394; pelobotryum, 394
Anacolia menziesii, 68+; menziesii var. baueri, 70
Anaptychia setifera, 394
Anastrophyllum minutum, 268
ANDERSON, L. E. *See* STONEBURNER, A., D. M. LANE & L. E. ANDERSON, 324
Andreaea, 174+; rothii, 174
Anomobryum, 263; prostratum, 263
Anomylia cuneifolia, 310
Anthoceros, 174+, 305+; donnellii, 309+; laevis, 8, 11; pearsonii, 75; phymatodes, 309; punctatus, 174
Anthocerotae, 305+
Anthocerotophyta, 305
anthraquinones, 52, 157
Anzia afromontana, 231
Aphanotropis, 42
Aplozia pendletonii, *75*
ARAVENA, R. & B. G. WARNER. Oxygen-18 composition of Sphagnum, and microenvironmental water relations, 445
Archilejeunea, 42+
Arctomia interfixa, 394
Arctoparmelia centrifuga, 1+, 378, 394; incurva, 394; separata, 394
Arthonia sp., 164, 392+; atomaria, 392+; **atomaria** comb. nov., 394; vinosa, 33+
arthothelin, 53
Arthrorhaphis alpina, 24, 394
ARUP, U. Caloplaca marina and C. rosei, two difficult species in North America, 148
ARUP, U. Caloplaca stantonii sp. nov. and its relationship to Caloplaca bolacina and other lobate to squamulose species in North America, 449
Asahinea chrysantha, 394
Ascochyta, 339; **candelariellicola** sp. nov., 338*; lichenoides, 339; santessonii, 339
Ascomycotina, 186
asemone, 53+
Aspicilia, 15+; aliena, 394; alphoplaca, 394; anseris, 394; caesiocinerea, 24, 394; candida, 394; cinerea, *139*, 394; disserpens, 394; gibbosa, 139; myrinii, 394; perradiata, 394; plicigera, 394; reptans, 330
Aspiromitus appalachianus, 305; laevis, 309+; lamellatus, 305; punctatus, 309
Asterella bolanderi, 74; californica, 75; drummondii, 10+
Astomiopsis amblyocalyx, 263
atranorin, 2+, 53, 79+, 377+, 430, 439+
Atrichum, 272, 324; angustatum, 324+; selwynii, 65; undulatum, 324+
Atwater, E. E., 74
Aulacomnium palustre, 409+
Aureolejeunea, 42

Bacidia bagliettoana, 394; herbarum, 394; subincompta, 24; trachona, 394
Baeomyces, 189+; arcuatus, 190; carneus, 24; heteromorphus, 190; roseus, 190; rufus, 394
Baeomycetaceae, 186+
Baker, M. S., 74
barbatic acid, 380
Barbilophozia floerkei, 268

[1] New scientific names and combinations are in boldface; page numbers of names in synonymy are in italic; taxa mentioned several times in an article are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

The Bryologist 95(4), 1992, pp. 492–501
Copyright © 1992 by the American Bryological and Lichenological Society, Inc.

Index to Volume 95 of THE BRYOLOGIST[1]

CYNTHIA M. GALLOWAY
Department of Biology, Texas A&I University, Kingsville, TX 78363

Abietinella abietina, 31
Abrothallus parmeliarum, 404; welwitzschii, 33+
Acarospora badiofusca, 33; fuscata, 139; glaucocarpa, 393; pelicypha, 393; schleicheri, 330; sinopica, 33, 393
Acrobolbus ciliatus, 309; rhizophyllus, 310
Adelanthus decipiens, 289
Adelolecia pilati, 393
Adelotheciaceae, 287
Adelothecium bogotense, 287+*
Adenostoma, 330; fasciculatum, 330
Aerobryidium subpiligerum, 289
Agrestia hispida, 33
Agyrophora lyngei, 393
Alaska, Calycularia crispula new to, 266; lichens of the Tuxedni Wilderness Area, 20
Alectoria nigricans, 24, 393; ochroleuca, 24, 393; vexillifera, 394
alectoronic acid, 2, 378+
alkaline phosphatase, 321+
Allantoparmelia almquistii, 394; alpicola, 394
ALLEN, B. [Review:] H. Crum. Liverworts and Hornworts of Southern Michigan, 374
ALLEN, B. [Review:] J.-P. Frahm. Campylopodes Centrali-Africanae, 371
ALLEN, B. [Review:] J.-P. Frahm. Campylopodes Exsiccatae Fasc. VII, Nr. 151–175, 371
ALLEN, B. [Review:] S. R. Gradstein. Bryophyta Neotropica Exsiccata. Fascicle VI, 371
ALLEN, B. *See* MAGILL, R. E. & B. ALLEN, 128, 363
ALLEN, B. Teratological axillary "hairs" in Didymodon tophaceus, 97
ALLEN, B. & R. E. MAGILL. Recent literature on mosses—152, 250; —154, 478
Allisoniaceae, 268
Aloina bifrons, 280
Alophosia, 272
Alsia californica, 69+
ALSTRUP, V. *See* ISBRAND, S. & V. ALSTRUP, 233
Amandinea, 436
Amblyolejeunea, 42
Amblystegium, 334
Ames, M. P., 73
Amygdalaria consentiens, 394; elegantior, 392+; panaeola, 394; pelobotryum, 394
Anacolia menziesii, 68+; menziesii var. baueri, 70
Anaptychia setifera, 394
Anastrophyllum minutum, 268
ANDERSON, L. E. *See* STONEBURNER, A., D. M. LANE & L. E. ANDERSON, 324
Andreaea, 174+; rothii, 174
Anomobryum, 263; prostratum, 263
Anomylia cuneifolia, 310
Anthoceros, 174+, 305+; donnellii, 309+; laevis, 8, 11; pearsonii, 75; phymatodes, 309; punctatus, 174
Anthocerotae, 305+
Anthocerotophyta, 305
anthraquinones, 52, 157
Anzia afromontana, 231
Aphanotropis, 42
Aplozia pendletonii, *75*
ARAVENA, R. & B. G. WARNER. Oxygen-18 composition of Sphagnum, and microenvironmental water relations, 445
Archilejeunea, 42+
Arctomia interfixa, 394
Arctoparmelia centrifuga, 1+, 378, 394; incurva, 394; separata, 394
Arthonia sp., 164, 392+; atomaria, 392+; **atomaria** comb. nov., 394; vinosa, 33+
arthothelin, 53
Arthrorhaphis alpina, 24, 394
ARUP, U. Caloplaca marina and C. rosei, two difficult species in North America, 148
ARUP, U. Caloplaca stantonii sp. nov. and its relationship to Caloplaca bolacina and other lobate to squamulose species in North America, 449
Asahinea chrysantha, 394
Ascochyta, 339; **candelariellicola** sp. nov., 338*; lichenoides, 339; santessonii, 339
Ascomycotina, 186
asemone, 53+
Aspicilia, 15+; aliena, 394; alphoplaca, 394; anseris, 394; caesiocinerea, 24, 394; candida, 394; cinerea, *139*, 394; disserpens, 394; gibbosa, 139; myrinii, 394; perradiata, 394; plicigera, 394; reptans, 330
Aspiromitus appalachianus, 305; laevis, 309+; lamellatus, 305; punctatus, 309
Asterella bolanderi, 74; californica, 75; drummondii, 10+
Astomiopsis amblyocalyx, 263
atranorin, 2+, 53, 79+, 377+, 430, 439+
Atrichum, 272, 324; angustatum, 324+; selwynii, 65; undulatum, 324+
Atwater, E. E., 74
Aulacomnium palustre, 409+
Aureolejeunea, 42

Bacidia bagliettoana, 394; herbarum, 394; subincompta, 24; trachona, 394
Baeomyces, 189+; arcuatus, 190; carneus, 24; heteromorphus, 190; roseus, 190; rufus, 394
Baeomycetaceae, 186+
Baker, M. S., 74
barbatic acid, 380
Barbilophozia floerkei, 268

[1] New scientific names and combinations are in boldface; page numbers of names in synonymy are in italic; taxa mentioned several times in an article are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

Barbula acuta, 66; bolanderi, *74*; calycina, 11+; cf. calycina, 10+; crinita, 9+; hornschuchiana, 10; luteola, 9; unguiculata, 11
BARCLAY-ESTRUP, P. *See* PFEIFFER, H. N. & P. BARCLAY-ESTRUP, 40
Bartram, E. B., 74
Bartramia ithyphylla, 31; menziesii, *75*
Bartramiopsis, 272; lescurii, 268
Bauer, J. A., 71
Bazzania ambigua, 144; tricrenata, 144, 269
BEDNAREK-OCHYRA, H. *See* OCHYRA, R., H. BEDNAREK-OCHYRA, T. PÓCS & M. R. CROSBY, 287
Bellemerea alpina, 394
Biatora vernalis, 24, 395
Bigelow, J. M., 69
Blasia, 268
Blepharostoma trichophyllum, 31, 146+, 268
Bolander, H. N., 70+
Brachelyma subulatum, 97
Brachydontium, 340+; curvisetum, 340; flexisetum, 340; intermedium, 340; **notorogenes** sp. nov. 340, 341*; olympicum, 340+; olsonii, 341+; trichodes, 340+
Brachymenium exile, 11; preissianum, 10
Brachythecium, 263; bolanderi, *74*; collinum, 75; rutabulum, 9+
BRASSARD, G. R. *See* HEDDERSON, T. A. & G. R. BRASSARD, 31, 443
Breutelia, 263
Brewer, W. H., 70+
Brigantiaea, 233
Britton, E. G. K., 74
BRODO, I. M. *See* TØNSBERG, T. & I. M. BRODO, 225
Brodoa oroarctica, 395
Bruchia bolanderi, 74
Bryaceae, 337
Bryocaulon divergens, 24, 395
Bryonora castanea, 395
Bryopteris, 42, 293
Bryoria chalybeiformis, 395; nitidula, 395; simplicior, 395; tenuis, 395; trichodes, 24, 395
Bryum, 263; altisetum, 10; arachnoideum, 293; argenteum, 9+, 324+; atwateriae, *74*; baueri, 74; billardieri, 11; bolanderi, *74*; campylothecium, 9+; canariense, 263; capillare, 11; dichotomum, 9; microchaeton, 263; miniatum, 74; pachytheca, 11+; pallescens, 263; pseudotorquescens, 293; sauteri, 9
BUCK, W. R. [Review:] P. L. Nimis & T. J. Crovells (eds.). Quantitative Approaches to Phytogeography, 372
BUCK, W. R. [Review:] A. R. Perry. Mosses and Liverworts of Woodland: A Guide to Some of the Commonest Species, 489
BUCK, W. R. A revision of Philophyllum (Hookeriales), 334
BUCK, W. R. & R. R. IRELAND. Symphyodon (Symphyodontaceae) in the Americas, 433
BUCK, W. R. & A. SCHÄFER-VERWIMP. Brachydontium notorogenes sp. nov. (Seligeriaceae) from southeastern Brazil, 340
Buellia, 79, 436; sect. Diplotomma, 85; arnoldii, 81; atrofuscata, 81; bahiana, 82; blasteniospora, *86*; callispora *84*+; coniops, 24; curtisii, 81; didyma, *86*; disciformis, 24, 81, 139; erubescens, 24; imshaugiana, 80+; nivalis, 395; papillata, 395; placodiomorpha, 80+; placodiomorphoides, 81; pleiotropa, *82*; procellarum, 87; pulverulenta 395; punctata, 330, 395; punctata fo. muscicola, 395; remensa, 81; restituta, *84*+; rinodinospora, 84+, *86*; subconnexa, *84*+
calcium oxalate crystals, 430
California, Plagiochila satoi new to, 144; Thamnolia vermicularis new to, 458
Callicostella, 336; lorifolia, *334*;
Caloplaca, 148+; ammiospila, 395; bolacina, 159, 449+*; celata, 395; cerina, 24, 395; cirrochroa, 395; crenularia, 395; flavovirescens, 457; fraudans, 395; holocarpa, 395; intermedia, 449; invadens, 395; irrubescens, 449; jungermanniae, 395; marina, 148+, 149*, subsp. americana, 453, var. flavogranulata, 157; microthallina, 157+; obliterans, 159; rosei, 148+, 149*, 453; saxifragarum, 392+, 395; scopularis, 158; sinapisperma, 395; sipeana, 449; squamosa group, 449+; **stantonii** sp. nov., 454*; stillicidiorum, 395; subsoluta, 449; tetraspora, 395; tiroliensis, 395
Calycularia 266; crispula, 266+, 267*; laxa, 266; radiculosa, 266
Calymperopsis **elimbatus** sp. nov., 284, 286*; **mandrakensis** sp. nov., 284, 285*
Calypogeja fissa, 65
Calyptothecium planifrons, 293
Campbell, D. H., 72
Camptothecium amesiae, *73*
Campylium stellatum, 409+
Campylopus, 263; introflexus, 9; pauper, 263; schimperi, 263; sharpii, 263
Campylostelium, 341
Candelariella, 339; aurella, 24, 338+, 395; placodizans, 395; terrigena, 330, 395; vitellina, 395
CANO, M. J., J. GUERRA & R. M. ROS. Crossidium seriatum (Pottiaceae, Musci) new to Europe, 280
Carbonea vorticosa, 395
carboxylic acid, 181
Cardotiella, 205+; secunda, 206+, 213*; schlotheimiaeformis, 206, *213*; subappendiculata, 214
Carter, A., 72
Catapyrenium cinereum, 34, 395
Catharinea, 272
Catinaria, 394; atomaria, 394
Catolechia wahlenbergii, 24
Cavernularia, 268; hultenii, 24, 226
Cephalolejeunea, 42
Cephalophysis leucospila, 396
Cephalozia bicuspidata, 268; pleniceps, 269
Cephaloziella divaricata, 66
Ceratodon purpureus, 10
Cetraria andrejevii, 24; commixta, 24; cucullata, 24, 396; delisei, 396; ericetorum, 24, 396; hepatizon, 2, 24; islandica, 396, subsp. crispiformis, 24, subsp. islandica, 24; kamczatica, 25; laevigata, 25, 396; nigricans, 396; nigricascens, 396; nivalis, 25, 396; tilesii, 396
Cetrelia chicitae, 139
Chandonanthus hirtellus, 289
Cheilolejeunea, 42
Chiloscyphus polyanthos var. rivularis, 66
Chlamydomonas, 174+
Chlorella, 174+
chloroatranorin, 430
2-chloroemodin, 53
2-chlorolichexanthone, 53
5-chlorolichexanthone, 52+
2-chloro-6-*O*-methylnorlichexanthone, 53
5-chloro-6-*O*-methylnorlichexanthone, 52+
7-chloro-6-*O*-methylnorlichexanthone, 52+
2-chlorononorlichexanthone, 52+
7-chlorononorlichexanthone, 52+
chloroxanthones, 52+
Cladia inflata, 189; moniliformis, 189
Cladina, 3, 459+; arbuscula, 25, 396; mitis, 25, 396; por-

tentosa, 4; portentosa subsp. pacifica, 459; rangiferina, 25, 396; stellaris, 25, 396
Cladonia, 330; amaurocraea, 25, 396; bacilliformis, 396; bellidiflora, 25, 396; cariosa, 396; carneola, 396; cenotea, 25, 396; chlorophaea, 25; coccifera, 25, 396; coniocraea, 25; cornuta, 25, 396; crispata, 25, 396; cyanipes, 396; decorticata, 25; deformis, 396; ecmocyna, 25, 396; fimbriata, 25, 396; furcata, 25, 139; gracilis, 396, subsp. gracilis, 25, 396, subsp. turbinata, 25, 396; grayi, 25; humilis, 25; macilenta, 458; macroptera, 25; macrophylla, 396; maxima, 25; metacorallifera, 25; phyllophora, 25; pleurota, 25; pocillum, 330, 396; polydactyla, 25; pyxidata, 396; scabriuscula, 25; southlandica, 189; squamosa, 396; stricta, 25, 396; subapodocarpa, 139; subfurcata, 25, 396; sulphurina, 25, 396; uncialis, 25, 396; verruculosa, 25
Claopodium bolanderi, 74, 268; crispifolium, 68; whippleanum, 76
Clathroporina, 233
Clauzadea monticola, 396
CLAYDEN, S. R. Chemical divergence of eastern North American and European populations of Arctoparmelia centrifuga and their sympatric usnic acid-deficient chemotypes, 1
Clypeococcum hypocenomyceae, 33+
Coccotrema, 227; cucurbitula, 227+
Coccotremataceae, 227
Coelocaulon aculeatum, 396
Coleochaete, 172+
Coleochaetium capense, *214*; secundum, *212*
Collema, 330; ceraniscum, 396; cristatum, 34; furfuraceum, 25; cf. nigrescens, 25; glebulentum, 397; subflaccidum, 25; tenax, 397; undulatum, 25
Collie, A., 69
confluentic acid, 225
connorstictic acid, 380
Conocephalum, 176*; conicum, 174
constictic acid, 230
constipatic acid, 380+
Coville, F. V., 70
CRANDALL-STOTLER, B. [Review:] H. D. Zinsmeister & R. Mues (eds.). Bryophytes. Their Chemistry and Chemical Taxonomy, 374
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 100, 235, 345, 461
CROOK, C. E. *See* ELIX, J. A. & C. E. CROOK, 52
CROSBY, M. *See* OCHYRA, R., H. B. BEDNAREK-OCHYRA, T. PÓCS & M. R. CROSBY, 287
Crossidium, 280; aberrans, 65, 280+; asirense, 280; crassinerve, 280; davidae, 280; deserti, 280; geheebii, 10, 280; rosei, 280; seriatum, 65, 280+*
Crossomitrium, 334
CRUM, H. Miscellaneous notes on the genus Sphagnum. 1–2, 273
CRUM, H. Miscellaneous notes on the genus Sphagnum. 3. New species from Brazil, 419
CRUM, H. *See* YANO, O. & H. CRUM, 343
Crustodiplodina, 339
Cryptomitrium tenerum, 66
cryptostictic acid, 230
Ctenidium, 334
Cyphelium tigillare, 397
Cyrtomnium hymenophylloides, 31

Dactylina arctica, 25, 397; beringica, 25; madreporiformis, 397; ramulosa, 397
Daltonia, 263
DAVISON, P. G. & D. K. SMITH. Calycularia crispula (Hepaticae) in the Aleutian Islands and Pacific Northwest of North America, 266
Dawsonia, 272
Degelia gayana, 231
dehydroconstipatic acid, 380
DELGADILLO M., C. Moss interchange: Bryofloristic similarities between Mexico and Colombia and the phytogeographical role of the Central American bridge, 261
DE LUNA, E. *See* MISHLER, B. D., P. H. THRALL, J. S. HOPPLE, JR., E. DE LUNA & R. VILGALYS, 172
4-*O*-demethylbarbatic acid, 380+
2-*O*-demethylpsoromic acid, 188
Dendrolejeunea, 42+; fruticosa, 43
Dendroligotrichum, 272
depsides, 52+
depsidones, 52+, 377+
Dermatocarpon, 15+, 233; miniatum, 15+, 25; reticulatum, 15+, 25; vagans, 15+, 34
Desmatodon convolutus, 11; guepinii, 280; latifolius, 31; leucostoma, 31
dibenzofurans, 52
2,4-dichlorolichexanthone, 53
2,7-dichlorolichexanthone, 53
4,5-dichlorolichexanthone, 53
2,7-dichloronorlichexanthone, 53
4,7-dichloronorlichexanthone, 53
5,7-dichloronorlichexanthone, 53
2,5-dichloro-6-*O*-methylnorlichexanthone, 53+
2,7-dichloro-3-*O*-methylnorlichexanthone, 52+
2,7-dichloro-6-*O*-methylnorlichexanthone, 52+
5,7-dichloro-3-*O*-methylnorlichexanthone, 53
4,5-dichloronorlichexanthone, 53
2,4-dichloronorlichexanthone, 52+
2,5-dichloronorlichexanthone, 53
4,5-dichloro-6-*O*-methylnorlichexanthone, 53
Dichodontium olympicum, 268
Dicnemoloma pallidum, 9
Didymodon acutus var. icmadophilus, 66; aronis, 280; rigidulus var. icmadophilus, 66; subtorquatus, 10+; tophaceus, 97+; 98*
diffractaic acid, 380
Dimelaena, 164, 437+; angelica, *439*; cf. australiensis, 53+, 55*; oreina, 436; weberi, 441
diploicin, 79+
Diplophyllum albicans, 268; plicatum, 146; taxifolium, 144, 268
Diploschistes muscorum, 18, 330+; scruposus, 25, 397
diploschistesic acid, 381
Diplotomma alboatrum, 397
Dirina paradoxa, 164
Dirinetum africanae, 164
Distichophyllum geheebii, *289*
Ditrichum heteromallum, 263; schimperi, 76
Douglas, D., 68
DOWNING, A. J. Distribution of bryophytes on limestones in eastern Australia, 5
Drepanocladus fluitans, 334; uncinatus, 263
Drew, K. M., 74
Drummondoideae, 205
Dumortiera hirsuta, 312

Eccremidium arcuatum, 11
Echinothecium glabrum, 234
Ectropothecium laevifolium, 434*; **novae-valesiae** comb. nov., 223
EGEA, J. M & P. TORRENTE. Two new species of Lecanactis from Baja California, 161
ELIX, J. A. & C. E. CROOK. The joint occurrence of chlo-

roxanthones in lichens, and a further thirteen new lichen xanthones, 52
ELIX, J. A. *See* KANTVILAS, G., J. A. ELIX & P. W. JAMES, 186
ELIX, J. A. & T. H. NASH, III. A synopsis of the lichen genus Psiloparmelia (Ascomycotina, Parmeliaceae), 377
emodin, 153+
Encalypta affinis subsp. affinis, 31+, subsp. macounii, 31+; alpina, 31; brevipes, 31+; brevicolla, 31+; ciliata, 31+; longicolla, 31+; procera, 31+; rhaptocarpa, 31+; vulgaris, 9+
Endocarpon pulvinatum, 34; tortuosum, 34
Entographa zonata, 225+
Entosthodon bolanderi, 74
Ephebe hispidula, 397; lanata, 397
Epipterygium tozeri, 65
Erioderma leylandii, 231; mollissimum, 226
erythrommone, 53
ESSLINGER, T. L. Recent literature on lichens—145, 108; —146, 241; —147, 351; —148, 466
esterase, 321+
Euopsis granatina, 397
Euosmolejeunea, 42
Eurhynchium brittoniae, *74*; confertum, 324; praelongum, 66, 76
Evernia mesomorpha, 167, 397; perfragilis, 397

Fabronia australis, 9; pusilla, 69
FAHSELT, D. *See* HAGEMAN, C. & D. FAHSELT, 316
fallacinal, 153+
Fimbriaria bolanderi, *74*; lescurii, *75*
Fissidens, 263; fontanus, 92*; leptocladus, 9; mexicanus, 92+; milobakeri, 74; tuerckheimii, 92+; pungens, 9+; taylori, 11; vittatus, 9+; wallisii, 263
Flavoparmelia, 378+; baltimorensis, 139, 378; caperata, 167+, 378; haysomii, 378
Flavopunctelia flaventior, 167; soredica, 167
Fosberg, F. R., 74
Fossombronia sp., 11
Frullania, 142; bolanderi, 66, 74; probosciphora, 9; squarrosula, 9+
Fulfordianthus, 42+; pterobryoides, 43; evansii, 43
Fulgensia bracteata, 397
fumarprotocetraric acid, 378+
Funaria glabra, 11; hygrometrica, 324+; producta, 11
Funariaceae, 337
Fuscidea mollis, 397
GALLOWAY, D. J. & L. A. WATSON-GANDY. Lepolichen coccophorus (Lichenized Ascomycotina, Coccotremataceae) in South America, 227
gangaleoidin, 430
Geothallus tuberosus, 73
Gigaspermum repens, 9+
GIGNAC, L. D. Niche structure, resource partitioning, and species interactions of mire bryophytes relative to climatic and ecological gradients in western Canada, 406
GILES EMORY, K. S. *See* THIERS, B. M. & K. S. GILES EMORY, 68
glucose-6-phosphate dehydrogenase, 321
glutamate dehydrogenase, 321
Glyphocarpa baueri var. baueri, *71*
Goniomitrium acuminatum, 11
GOWARD, T. & G. THOR. Notes on the lichens and allied fungi of British Columbia. I, 33
GRADSTEIN, S. R. The genera Thysananthus, Dendrolejeunea, and Fulfordianthus gen. nov. (studies on Lejeuneaceae subfamily Ptychanthoideae XXI), 42
Grimmia affinis, 31; laevigata, 10+; moxleyi, 75; olympica, 341; pulvinata, 9+; rivularis, 263
griseoxanthone C, 52+
GUERRA, J. *See* CANO, M. J., J. GUERRA & R. M. ROS, 280
Gyalecta foveolaris, 397; geoica, 397
Gymnomitrion concinnatum, 268; laceratum, 269; pacificum, 269
Gymnoschoenus sphaerocephalus, 186+
Gymnostomum aeruginosum, 9+, 65
gyrophoric acid, 225, 437
Gyrothyra underwoodiana, 76

Haematomma lapponicum, 25
Hafellia, 79+; **bahiana** comb. nov., 82*, var. bahiana, 79+, var. **pleiotropa** comb. nov., 82; **bispora** sp. nov., 83*; **callispora** comb. nov. 84*; **dissa** comb. nov., 87; **fosteri** comb. nov., 85; leptoclinoides, 81+; parastata, 79+*, 86; **procellarum** comb. nov., 87
HAGEMAN, C. & D. FAHSELT. Geographical distribution of enzyme polymorphisms in the lichen Umbilicaria mammulata, 316
Halecania alpivaga, 397
Harpalejeunea ovata, 309
Harthill, M., 73
HAWKSWORTH, D. L. & K. KALB. A new species of Ascochyta on Candelariella aurella from Baja California, Mexico, 338
HEDDERSON, T. A. & G. R. BRASSARD. Encalypta affinis subsp. macounii and E. brevipes new to eastern North America from the Torngat, Northern Labrador, Canada, 31
HEDDERSON, T. A & G. R. BRASSARD. Tetrodontium repandum and Seligeria diversifolia discovered in arctic Canada, 443
Hedwigia ciliata, 9
Hedwigidium integrifolium, 9
Helocarpon crassipes, 397
Helotiales, 186
Herbertus aduncus, 268, 309
Heterodermia obscurata, 167; speciosa, 167
Hillebrand, W., 75
Homalothecium pinnatifidum, 73
HONG, W. S. Plagiochila in western North America, 142
Hookeria bogotensis, *287*+, 334; lorifolia, *334*; tenuifolia, *334*
Hookeriaceae, 287
HOPPLE, J. S., JR. *See* MISHLER, B. D., P. H. THRALL, J. S. HOPPLE, JR., E. DE LUNA & R. VILGALYS, 172
Howe, M. A., 72
Hylocomium splendens, 409+
Hymenelia lacustris, 397
Hyperphyscia, 436
Hypnum, 334+; bigelovii, *74*; bolanderi, *74*; brewerianum, 70, *74*; cupressiforme, 9; hillebrandii, *75*; revolutum, 31; royae, *76*; whippleanum, 69, *76*
Hypocenomyce leucococca, 33+; scalaris, 397
hypoconstictic acid, 380
Hypogymnia, 380; austerodes, 397; enteromorpha, 25; inactiva, 25; physodes, 25, 38+, 167+, 397; pseudophysodes, 33+; subobscura, 397; vittata, 25, 397
hyposalazinic acid, 380
hypostictic acid, 380

Icmadophila, 190; ericetorum, 190, 397
Imshaugia aleurites, 139, 167
Ionaspis epulotica, 398; melanocarpa, 398; schismatopsis, 398; suaveolens, 398
IRELAND, R. R. Studies of the genus Plagiothecium in Australasia, 221
IRELAND, R. R. *See* BUCK, W. R. & R. R. IRELAND, 433

ISBRAND, S. & V. ALSTRUP. Opegrapha sphaerophoricola sp. nov., 233
isoarthothelin, 53
isocitrate dehydrogenase, 321
isolichenan, 377+
isonotatic acid, 380
Isopterygium machrisianum, *433*; tenerifolium, *334*
Isothecium cristatum, 75; howeii, 75; myosuroides, 68
isousnic acid, 380
isozymes, 316

JAMES, P. W. *See* KANTVILAS, G., J. A. ELIX & P. W. JAMES, 186
Japewia tornoensis, 398
Jubula, 309
Jungermannia bolanderi, 74; pendletonii, 75

KALB, K. *See* HAWKSWORTH, D. L. & K. KALB, 338
Kantia trichomanis, 66
KANTVILAS, G., J. A. ELIX & P. W. JAMES. Siphulella, a new lichen genus from southwest Tasmania, 186
Kiaeria starkei, 31
Kindbergia praelonga, 74
Knightiella, 190; splachnirima, 189+
Koch, L. F., 72
Kurzia makinoana, 268
Kymatolejeunea, 42

LANE, D. M. *See* STONEBURNER, A., D. M. LANE & L. E. ANDERSON, 324
Lasallia, 380
LAWREY, J. D. Naturally and randomly-assembled lichen communities compared using the species-area curve, 137
Lecanactis dimelaenoides, 161+; nashii, 161+; werneri, 164; zahlbruchneri, 164
Lecania arctica, 398; erysibe, 398
Lecanora, 52+, 54*, 227, 233, 377, 430+; allophana, 398; argentea, 430+; argopholis, 398; atrosulphurea, 398; behringii, 52+; broccha, 52+; campestris, 139; cenisia, 398; cinererofusca, 25; circumborealis, 25, 398; coccophora, *228*+; contractula, 52+, 398; depressa, 432; dispersa, 398; epanora, 25; epibryon, 398; fuliginosa, 430+; gangaleoides, 430; garovaglii, 18; grantii, 85; hagenii, 398; hybocarpa, 25; hypoptoides, 398, cf. impudens, 25; intricata, 25, 398; luteovernalis, 398; marginata, 398; marina, 155; meiococca, 54; milvina, 436; muralis, 330; murorum var. lobulata, 157; nordenskioeldii, 398; pacifica, 25; pallida, 25; phaedrophthalma, 18; piniperda, 26; polytropa, 26, 398; populicola, 52+; pulicaris, 26; reagens, 33+; rupicola, 26, 398; salina, 52+; sophodes, 436; subfusca, 430+; sulphurata, 54+; sulphurescens, 431+; symmicta, 26, 398; torrida, 398; turfacea, 436; varia, 26, 398; vincentina, 430+; vinetorum, 52+; vorax, 52+; zosterae, 398
Lecidea, 377; albohyalina, 26; alpestris, 398; atrobrunnea, 398; auriculata, 26, 398; callispora, *84*; cinnabarina, 26, cf. elabens, 26; circumnigrata, 398; confluens, 398; cuprea, 398; diapensiae, 398; dissa, *87*; ementiens, 398; hypnorum, 398; hypocrita, 26, 398; lactea, 398; lapicida, 26, 398, fo. ochracea, 26, 398; limosa, 398; nylanderi, 398; parastata, *86*; picea, 398; plebeja, 399; pullata, 33+; ramulosa, 399; restituta, *84*; silacea, 35; subconnexa, *84*; tessellata, 26, 399; umbonata, 399
Lecidella, 52+; alaiensis, 399; elaeochroma, 399; euphorea, 26, 399; inamoena, *399*; meiococca, 52+; potaviana, 399; vorax, 54; wulfenii, 399
Lecidoma demissum, 399
Leciographa, 233
Leciophysma finmarkicum, 399
Leioderma pycnophorum, 231
Leiodon, 272
Leiomitrium capense, 214
Lejeunea ulicina subsp. ulicina, 309
Lejeuneaceae, 289
Lempholemma cladodes, 399
leoidin, 430
Lepidopilum, 263
Lepidozia reptans, 146
Lepolichen, 227+; coccophorus, 227+*; granulatus, 227
Lepraria, 26; zonata, 139
Leptodon smithii, 9+
Leptodontium, 263; proliferum, 263
Leptogium lichenoides, 399; minutissimum, 399; rivale, 33+; saturninum, 26; sinuatum, 330
Leptomitaceae, 380
Leptomitales, 380
Leptoscyphus anomalus, 309
Leptotrichum schimperi, *76*
LESICA, P. *See* MCCUNE, B. & P. LESICA, 296
Leskea angustata, 263
Lesquereux, C. L., 75
Leucolepis acanthoneuron, 68
Leucomiaceae, 335+
Leucomium, 337; connexum, 336*; strumosum, 336*
Levierella perserrata, 293
LEWINSKY, J. Orthotrichum truncato-dentatum, a species known only from extinct habitats, 218
Lichen hepatizon, 403
lichenan, 377; Cetraria-type, 378+; Xanthoparmelia-type, 378+
lichexanthone derivatives, 54*
Lobaria hallii, 26; linita, 26; pulmonaria, 26, 217
Lobariaceae, 380
LONG, R. P. *See* SHOWMAN, R. E. & R. P. LONG, 166
Lopadium coralloideum, 399; pezizoideum, 399
Lophocolea semiteres, 9
Lophozia cf. alpestris, 268; collaris, 146; heterocolpa, 147
LUMBSCH, H. T. Lecanora argentea Oxner & Volkova, the correct name for L. fuliginosa Brodo, and comments on related species, 430
Lunularia cruciata, 10+
Lyellia, 272

Macrocoma, 205+; filiforme, 214; frigida, 263; lycopodioides, 205+, 207*; pulchella, 205+, 207*; tenue, 205+, subsp. sullivantii, 208, subsp. tenue, 206, 207*
Macromitrioideae, 205+
Macromitrium, 205+, 263; subgen. Macrocoma, 214; sect. Macrocoma, 214; confusum, *214*; dawsoniaemitrium, *214*; dregei, *214*; ferrugineum, 214; lebomboense, 205+, 209*, 214; levatum, 205+, 209*, 289; ligulare, 210; lycopodioides, *214*; macropelma, 205+; mannii, *208*; microphyllum, 214; pallidum, 208; pulchellum, *214*; richardii, 205+; schlotheimiaeformis, *213*; secundum, 205+, *212*; serpens, 205+, 209*; tenue, *214*, var. dregei, 214, var. lycopodioides, *214*, fo. microphyllum, *214*; tristratosum, 205+
Madotheca bolanderi, *74*; douglasii, *69*+
MAGILL, R. E. *See* ALLEN, B. & R. E. MAGILL, 250, 478
MAGILL, R. E. & B. ALLEN. Recent literature on mosses—151, 128; —153, 363
Malaysia, Syrrhopodon stonae new to, 88
malonprotocetraric acid, 386
mannitol dehydrogenase, 319
Marchantia, 173+; polymorpha, 66, 173+
Marsupella bolanderi, 66+
Maschalocarpus ecklonii, 214

Massalongia carnosa, 399
Mastigolejeunea, 42
MATZER, M. *See* MAYRHOFER, H., J. W. SHEARD & M. MATZER, 436
MAYRHOFER, H., J. W. SHEARD & M. MATZER. Mobergia (Physciaceae, Lichenized Ascomycetes), a new genus endemic to western North America, 436
McClatchie, A. J., 71
MCCUNE, B. & P. LESICA. The trade-off between species capture and quantitative accuracy in ecological inventory of lichens and bryophytes in forests in Montana, 296
MCCUNE, B. & R. ROSENTRETER. Texosporium sancti-jacobi, a rare western North American lichen, 329
MCCUNE, B. *See* ROSENTRETER, R. & B. MCCUNE, 15
McFadden, F. M., 71
Megaceros, 305+; aenigmaticus, 305+*; tosanum, 314
Megaspora verrucosa, 399
Meiotrichum gen. & stat. nov., 270; **lyallii** comb. nov., 270
Melanelia disjuncta, 399; elegantula, 399; exasperatula, 26; granulosa, 399; incolorata, 26, 399; infumata, 399; olivacea, 26, 399; panniformis, 399; septentrionalis, 26; sorediata, 399; stygia, 26, 399; subaurifera, 167
Menegazzia, 380; hollermayeri, 231; valdiviensis, 231
menegazziaic acid, 230
Menzies, A., 68
MERRILL, G. L. SMITH. Notes on North American Polytrichaceae: Polytrichastrum, 270
Metaneckera menziesii, 68
6-*O*-methylarthothelin, 53
3-*O*-methylasemone, 53
6-*O*-methylasemone, 53
methyl 2,5-dichloronorpsoromate, 188
methyl 2,5-dichloropsoromate, 188
methylhypoprotocetraric acid, 380
2-*O*-methylmicrophyllinic acid, 225
3-*O*-methylnorlichexanthone, 52+
3-*O*-methylnorlichexanthone derivatives, 54*
6-*O*-methylnorlichexanthone derivatives, 54*
3-*O*-methylthiophanic acid, 52+
6-*O*-methylthiophanic acid, 52+
Metzgeria, 174+; conjugata, 268; furcata, 174, 309, var. ulvula, 309, var. setigera, 309
Metzgeriales, 268
Micarea assimilata, 26, 399; austroternaria, 189; crassipes, 26; denigrata, 26; isabellina, 52+, 189; incrassata, 399; melaena, 399; ternaria, 399
Microlychnus epicorticis, 35+
Miriquidica leucophaea, 399; lulensis, 399
MISHLER, B. D., P. H. THRALL, J. S. HOPPLE, JR., E. DE LUNA & R. VILGALYS. A molecular approach to the phylogeny of bryophytes: Cladistic analysis of chloroplast-encoded 16S and 23S ribosomal RNA genes, 172
Mitthyridium jungquilianum, 91; papuanum, 88+; subluteum, 88+
Mniadelphus geheebii, *289*
Mnium, 309
Mobergia gen. nov., 438; **angelica** comb. nov. 439*; **calculiformis** comb. nov., 441*
Moerckia, 266
MOHAMED, J. & W. D. REESE. Syrrhopodon stonae, new to Malaya, and notes on other Malayan Calymperaceae, 88
Moxley, G. L., 75
Multiclavula corynoides, 35; sharpii, 35
Mycobilimbia berengeriana, 399
Mycoblastus alpinus, 26; sanguinarius, 26, 399
NASH, T. H., III. *See* ELIX, J. A. & T. H. NASH, III, 377
Neckera californica, *69*; douglasii, 68+; pennata, 97+, fo. pterantha, 97+, var. tenera, 97+
Neorutenbergia, 287
Nephroma antarcticum, 231; arcticum, 26, 400; bellum, 26; expallidum, 400; laevigatum, 226; parile, 26; resupinatum, 217
norgangaleoidin, 430
norlichexanthone, 52+; derivatives, 54*
Norris, D., 73
norstictic acid, 79+, 189, 230, 436
notatic acid, 380
Notoligotrichum, 272

obtusatic acid, 380
Ochrolechia androgyna, 400; arborea, 26; frigida, 26, 400; grimmiae, 400; inaequatula, 400; laevigata, 26; oregonensis, 26; szatalaensis, 26; upsaliensis, 400; yasudae, 139
OCHYRA, R., H. BEDNAREK-OCHYRA, T. PÓCS & M. CROSBY. The moss Adelothecium bogotense in continental Africa, with a review of its world range, 287
Octodiceras, 92+
Oligotrichum, 272
Omphalanthus, 42
Omphalodiscus decussatus, 400; virginis, 400
Opegrapha, 233+; gyrocarpa, 225+; **sphaerophoricola** sp. nov., 233*
Opegraphaceae, 233
Ophioparma lapponica, 400
β-orcinol depsidone, 186+, 377+
Oreoweisia lechleri, 263
Orphniospora moriopsis, 400
orthophosphoric acid, 53
Orthotrichaceae, 205
Orthotrichoideae, 205
Orthotrichum, 205+; alpestre, 31+; araucarieti, 218+; cupulatum var. cupulatum, 9+; ferrugineum, 214; hallii, 65; kingianum, 75; lyellii, 69; microphyllum, 215; serpens, 215; tenue, 215; trachymitrium, 263; truncato-dentatum, 218+, 219*
oxalate crystals, 438+

Pallavicinia lyellii, 268
Pallaviciniaceae, 268
Pannaria confusa, 217; hookeri, 400; pezizoides, 26, 400; praetermissa, 400
Pannoparmelia anzioides, 231
Paraleucobryum enerve, 263+
parietin, 153+
Parmelia 233, 377; aleuritica, 1+; aurulenta, 167; bouly de lesdainii, 386; centrifuga β dealbata, *1*, fo. aleuritica, *1*, fo. dealbata, 1, var. dealbata, 1+; coccophora, 227+; distincta, 378, *384*, 385*, fo. subcongruens, *384*; fraudans, 400, 404; omphalodes, 26, 139, 400; saxatilis, 26, 400; squarrosa, 167; subcongruens, *384*+; sulcata, 26, 167+, 217, 400
Parmeliaceae, 377, 380
Parmeliopsis aleurites, 400; ambigua, 3+, 400; hyperopta, 3+, 26, 400
Parmotrema crinitum, 139; stuppeum, 167
Pearson, W. H., 75
Pellia, 266+; endiviifolia, 268
Pelliaceae, 268
Peltigera aphthosa, 26, 400; canina, 26, 167, 400; collina, 26; didactyla, 26, 400; horizontalis, 26; lepidophora, 400; leucophlebia, 400; malacea, 26, 400; membranacea, 26; neckeri, 26; neopolydactyla, 27; polydactyla,

27, 400; rufescens, 27, 400; scabrosa, 27, 400; venosa, 27, 400
Pendleton, G. M., 75
Perforaria granulata, 227
Pertusaria, 139, 227, 233; alpina, 27; carneopallida, 27; cicatricosa, 54, 55*; coccophora, *228*; coriacea, 400; cucurbitula, 277; dactylina, 27, 400; gymnospora, 189; leucostoma, 27, cf. macounii, 27; octomela, 400; oculata, 27, 400; ophthalmiza, 27; panyrga, 27, 401; pycnothelia var. A, 52+, 55*; sommerfeltii, 27; subdactylina, 27; subobducens, 401; thamnoplaca, 227; trochiscea, 401; xanthostoma, 27
pertusaric acid, 380
Pfeiffer, H. N. & P. Barclay-Estrup. The use of a single lichen species, Hypogymnia physodes, as an indicator of air quality in northwestern Ontario, 38
Phaeoceros laevis, 66
Phaeophyscia, 436; adiastola, 35; endococcinea, 401; hirsuta, 33+; hirtella, 27+; nigricans, 35; orbicularis, 27; pusilloides, 167; rubropulchra, 167; sciastra, 27, 339, 401
Phaeorrhiza, 233; nimbosa, 401; sareptana, 330
Phaeospora parasitica, 404
Phascum cuspidatum, 66; hyalinotrichum, 66; vlassovii, 280
Philonotis, 263; marchica, 263
Philophyllum, 334+; bromeliae, *334*; bromeliophilum, *334*; lorifolium, *334*; tenuifolium, *344*+, 336*
Phlyctis argena, 217
6-phosphogluconate dehydrogenase, 321
Phylliscum demangeoni, 401
Phyllogonium, 293
Phyllopsora corallina var. ochroxantha, 188
Phyllopyreniaceae, 227
Phyllopyrenieae, 227
Physcia, 436; adscendens, 27, 401; aipolia, 27, 401; caesia, 27, 401; dimidiata, 33+; dubia, 27; millegrana, 167, semipinnata, 35; stellaris, 167; tenella, 27
Physciaceae, 81, 436+
Physcomitrella, 173+; patens, 172
Physconia detersa, 27; distorta, 401; gelida, 27; grisea, 167; muscigena, 27, 401; perisidiosa, 27
Pilophorus cereolus, 35
Placidiopsis cervinula, 401
Placodium bolacinum, *450*
Placopsis gelida, 401
Placynthium asperellum, 401; nigrum, 401; stenophyllum var. isidiatum, 36
Plagiobryum zierii, 32
Plagiochasma rupestre, 11+
Plagiochila, 142+; arctica var. intermedia, *147*; asplenioides, 142; asplenioides subsp. arctica, 142+, 144*; asplenioides subsp. asplenioides, 142, 143*; asplenioides subsp. porelloides, 142, 143*; asplenioides subsp. porelloides, 142, 143*; asplenioides subsp. porelloides fo. subarctica, 142+, 144*; asplenioides subsp. porelloides var. grandiretis, *147*; austini, 309; barteri, 289; corniculata 309; major, 142; satoi, 142; satoi var. **magnum** var. nov., 144*; satoi var. satoi, 142+, 143*; schofieldiana, 142, 143*; semidecurrens, 142; semidecurrens var. alaskana, 142+, 143*; semidecurrens var. semidecurrens, 142+, 143*; sullivantii, 309; yokogurensis, 309
Plagiothecium, 221+; denticulatum, 221; laetum, 268; lucidum, 221+; novae-seelandiae, 221+*
Platismatia glauca, 27, 401; lacunosa, 27; norvegica, 226; tuckermanii, 167
Pleuridium acuminatum, 74; bolanderi, *74*; californicum, 66+; nervosum, 10; subulatum, 66
Pleurozium schreberi, 39, 268, 409+, 410*
Pócs, T. *See* Ochyra, R., H. Bednarek-Ochyra, T. Pócs & M. Crosby, 287
Pogonatum, 270+; alpinum, 268; contortum, 68; sphaerothecium, *270*
Pohlia filiformis, 268; longicollis, 74
Polyblastia bryophila, 401; cruenta, 401; cupularis, 401; hyperborea var. hyperborea, 401, var. macrospora, 401; melaspora, 392+, 401; obsoleta, 401; sendtneri, 401; theleodes, 401
polygalacturonic acid, 181
polyketides, 52
polymerase chain reaction, 173*
Polysporina simplex, 401
Polytrichaceae, 270+, 324
Polytrichadelphus, 271+; lyallii, *271*
Polytrichastrum, 270+; sect. **Aporotheca** comb. nov., 270; sect. Meiotrichum, *271*; sect. Polytrichastrum, 270; sect. Sexangularia, *270*+; alpiniforme, *270*; alpinum, 270+, var. fragile, 270, var. septentrionale, 270; **appalachianum** comb. nov., 270; emodi, 270+; formosum, 271; longisetum, 271+; lyallii, 271; ohioense, 271+; papillatum, 270+; pallidisetum, 271; sexangulare, 270+; sexangulare var. **vulcanicum** comb. nov., 270; tenellum, 270+; torquatum, 271; xanthopilum, 271
Polytrichum, 68, 172+, 270+; sect. Aporotheca, *271*; angustifolium, 271; appalachianum, *271*; commune, 174, 263, 271; ericoides, 271; formosum, 271; hyperboreum, 271; juniperinum, 271, 324; longisetum, 68; piliferum, 271, 324; sexangulare, *270*; sphaerothecium, 270; strictum, 271; swartzii, 271
polyuronic acid, 181
Porella, 172+; bolanderi, 66, 74; navicularis, 68+; pinnata, 174
Porina granulata, 227+
Porotrichum bigelovii, 74
Porpidia albocaerulescens, 139; flavocaerulescens, 401; macrocarpa, 27, 401; speirea, 401; thomsonii, 401; zeoroides, 401
Pottia, 10; fosbergii, *75*; starkeana, 75
Pottiaceae sp. b, 10
Protoblastenia rupestris, 401; terricola, 402
protocetraric acid, 380+
protoconstipatic acid, 380
Protoparmelia badia, 402
Pseudephebe minuscula, 27, 402; pubescens, 27, 402
Pseudocrossidium revolutum, 280
Pseudocyphellaria berberina, 231; flavicans, 231; hirsuta, 231; nitida, 231; pluvialis, 231
Pseudoleskeopsis imbricata, 9+
Pseudotaxiphyllum elegans, 68; machrisianum, *433*
Psiloparmelia, 377+; sect. Xanthoparmelia subsect. Endocoerulea, *382*+; aleuritica, 378; arhizinosa, 379+, 383*; centrifuga, 378; **denotata** sp. nov., 382*; **dichotoma** sp. nov., 384; **diffractaica** sp. nov., 384*; distincta, 378+, 385*; **flavobrunnea** comb. nov., 386*; **hypostictica** sp. nov., 387*; incurva, 378; **norstictica** sp. nov., 387*; **pustulata** sp. nov., 388*; **salazinica** sp. nov., 388*; separata, 378; **sorediosa** sp. nov., 388*; subcentrifuga, 378; **subcrustosa** sp. nov., 390*
Psilopilum, 272+
Psora, 330; decipiens, 402; rubiformis, 402
Psoroma hispidulum, 231; hypnorum, 27, 402; microphyllizans, 231
psoromic acid, 186+
Pterogonium gracile, 69
Pterygodon, 272
Pterygoneurum koslovii, 280
Ptilidium ciliare, 146

Ptychanthus, 42
Ptychomitriaceae, 341
Ptychomitrium australe, 9; gardneri, 75
Punctelia rudecta, 167+; subrudecta, 27, 167+
PURSELL, R. A. The occurrence of fertile Fissidens fontanus in Mexico, 92
Pyrenocollema elegans, 157
Pyrenopsis pulvinata, 402
Pyxine sorediata, 167

Racomitrium heterostichum, 65; lamprocarpum, 293; microcarpon, 31
Radula bolanderi, 74; voluta, 309
Ramalina farinacea, 27; intermedia, 139; menziesii, 192+*; roesleri, 27, 402; scoparia, 402
ravenelin, 53
REDFEARN, P. L., JR. & R. E. MAGILL. Three species of Fissidens new to mainland China, and a second collection of F. brevinervis (Musci: Fissidentaceae), 171
REESE, W. D. Syrrhopodon cyrtacanthos, a new species of moss from Australia, 94
REESE, W. D. *See* MOHAMED, H. & W. D. REESE, 88
Reinkella parishii, 164
Rhipidiaceae, 380
Rhizocarpon anseris, 392+, 402; atroflavescens, 402; badioatrum, 402; chioneum, 402, 404; copelandii, 402; crystalligenum, 402; distinctum, 402; eupetraeoides, 27, 402; eupetraeum, 27, 402; ferax, 402; geminatum, 27, 402; geographicum, 27, 402; grande, 27, 402; hochstetteri, 402; inarense, 27, 402; polycarpum, 27, 402; rittokense, 402; superficiale, 402; umbilicatum, 402
Rhizomnium magnifolium, 268
Rhizoplaca, 15+; chrysoleuca, 27, 139
Rhodobryum grandifolium, 263+
Rhynchostegiopsis, 334+
Rhynchostegium, 334+
Riccardia chamedryfolia, 66+, 305+; major, *66*; multifida, 305+
Riccia, 330; bifurca, 11+; californica, 66; campbelliana, 74; cavernosa, 66; limbata, 11+; sorocarpa, 66
Rimularia insularis, 402
Rinodina, 436+; sect. Beltraminia, 437; albana, 437; angelica, 436+, 437*, *439*; archaea, 27, 402; arnoldii, 436; atrocinerea, 437*; badiella, 436; bolodes, *436*+; calcigena, 402; calculiformis, 436+, 437*, *441*; confragrosa, 437*; conradii, 436; corticola, 437; dirinoides, *436*+; dissa, *87*; efflorescens, 36; insperata, 81; milvina, 402, 436; mniaraea, 402; oxydata, 81; petermannii, 436, 437*; platyloba, *436*+; pyreniospora, *436*; pyrina, 402; roscida, 402; septentrionalis, 27; **sheardii** sp. nov., 216*; sophodes, 436, 437*; subminutae, 216; subsophodes, 402; trachytica, 81; turfacea, 403, 436
Roccella babingtonii, 164; fimbriata, 164
roccellic acid, 431
ROS, R. M. *See* CANO, M. J., J. GUERRA & R. M. ROS, 280
ROSENTRETER, R. *See* MCCUNE, B. & R. ROSENTRETER, 329
ROSENTRETER, R. & B. MCCUNE. Vagrant Dermatocarpon in western North America, 15
Roy, J. D. G., 76
Rutenbergia, 287
Rutenbergiaceae, 287
RYAN, B. D. [Review:] P. M. McCarthy. Bibliography of Australian Lichens: 1807–1991, 488

Sagenotortula quitoensis, 263
Sagiolechia rhexoblephara, 27, 403
salazinic acid, 53, 381
sampling methods, 296+; belt-transect method, 297+; whole plant method, 297+
Sandeanthus, 43
Sandeothallaceae, 266
Sandeothallus, 266
SANDERS, W. B. Comparative in situ studies of thallus net development in morphologically distinct populations of the lichen Ramalina menziesii, 192
Sarcogyne, 139
Sarcoscyphus bolanderi, *74*
Scapania americana, 144; bolanderi, 74, 146; parvifolia, 147; umbrosa, 146
Schaereria tenebrosa, 403
SCHÄFER-VERWIMP, A. *See* BUCK, W. R & A. SCHÄFER-VERWIMP, 341
Schimperobryum, 336
Schistidium apocarpum, 9
schizopeltic acid, 225
Schlotheimia, 205+; exrugulosa, *212*; ferruginea, 205+, 211*, 289; grevilleana, 205+; percuspidata, 205+, 211*; pulchella, 214; rufoaeruginosa, 205+, 211*; rufoglauca, 214; rufopallens, 211*, 212+; subventrosa, *212*; torquata, 210; ventrosa, 205+
SCHUSTER, R. M. On Megaceros aenigmaticus, 305
Sclerophyton sp., 164
Scoliosporum chlorococcum, 27
Seligeria diversifolia, 443+*; pusilla, 444
Sematophyllum, 263
Sewardiella, 268
SHEARD, J. W. The lichenized ascomycete genus Hafellia in North America, 79
SHEARD, J. W. *See* MAYHOFER, H., J. W. SHEARD & M. MATZER, 436
SHOWMAN, R. E. & R. P. LONG. Lichen studies along a wet sulfate deposition gradient in Pennsylvania, 166
Siphula decumbens, 189; jamesii, 189
Siphulella gen. nov., 186; **coralloidea** sp. nov., 186*
siphulellic acid, 188
skyrin, 430
SMITH, D. K. *See* DAVISON, P. G. & D. K. SMITH, 266
Solorina bispora, 403; crocea, 27, 403; spongiosa, 403
Spain, Crossidium seriatum new to, 280
Sphaerocarpales, 73
Sphaerocarpos, 172+; cristatus, 73; drewei, 74; texanum, 174
Sphaerophorus, 234; fragilis, 27, 403; globosus, 27, 233, 403; tener, 231; tuckermanii, 27
Sphagnum, 172+, 263, 268, 324, 343, 419; sect. Acutifolia, 276; sect. Buchanania, 276; sect. Cuspidata, 276; sect. Hemitheca, 276; sect. Isocladus, 274+; sect. Inretorta, 276; sect. Insulosa, 276; sect. Litophloea, 275+; sect. Polyclada, 276; sect. Rigida, 276; sect. Sphagnum, 276; sect. Squarrosa, 276; sect. Subsecunda, 276; subg. Acisphagnum, 276; subg. Homophylla, 276; subg. Inophloea, 276; subg. Isocladus, 274+; subg. Malacosphagnum, 276; subg. Rigida, 276; subg. Sphagnum, 276; subsect. Sericea, 275+; subsect. Cuspidata, 276; subsect. Subsecunda, 275+; **acutirameum** sp. nov., 423*; angustifolium, 406+; austinii, 406+; **bourbonense** sp. nov., 276*; brasiliense, 419; brevirameum, 419; **buckianum** sp. nov., 419*; capense, 278; capillifolium, 406+; 445+; compactum, 174; contortum, 276; cuspidatum, 447; davidii, 277+; **divisum** sp. nov., 424*; efibrillosum, 274+, 275*; **exquisitum** sp. nov., 420*; fallax, 276, 445+; fuscum, 409+, 445+; **geraisense** sp. nov., 424*; **homophyllum** sp. nov., 425*; inundatum, 276; irwinii, 419, 420*; itacolumitis, 419; lindbergii, 409+; longistolo, 343; macrophyllum, 274+; magel-

lanicum, 409+, 445+; molle, 447; novae-caledoniae, 275+; obovatum, 278; **obliquefibrosum** sp. nov., 426*; pacificum, 409+; papillosum, 406+, 445+; ramulinum, 423; recurvum, 276; rubellum, 406+; rutenbergii, 278; sericeum, 275+; splendens, 275; squarrosum, 181+*; submedium, 420+; subtile, 324+; **tabuleirense** sp. nov., 343, 344*; tenellum, 406+; warnstorfii, 409+
Sphenolobopsis pearsonii, 310
Spilonema revertens, 403
Sporastatia polyspora, 403; testudinea, 403
Spruceanthus, 42; polymorphus, 42
Squamidium brasiliense, 293
STARK, L. R. & A. T. WHITTEMORE. Additions to the bryoflora of southern California, 65
Staurothele, 233; drummondii, 27, 403; fissa, 403
Steere, W. C., 73
Stenorhipis madagascariensis, 289
Stereocaulon, 189, 233; alpinum, 27, 403; arenarium, 403; botryosum, 27, 403; cf. condensatum, 27; glareosum, 403; grande, 27; intermedium, 27; paschale, 27, 403; rivulorum, 27, 403; saxatile, 27; vesuvianum, 403
Sticta caulescens, 231; hypochra, 231
stictic acid, 189, 230
Stigmidium schaereri, 27
STONEBURNER, A., D. M. LANE & L. E. ANDERSON. Spore dispersal distances in Atrichum angustatum (Polytrichaceae), 324
STOTLER, R. [Review:] M. O. Hill, C. D. Preston & A. J. E. Smith (eds.). Atlas of the Bryophytes of Britain and Ireland. Volume 1. Liverworts (Hepaticae and Anthocerotae), 373
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—127, 100; —128, 235; —129, 345; —130, 461
Strepsilejeunea, 42
succinoprotocetraric acid, 386
superoxide dismutase, 321
Sutliffe, D., 72
Symbiezidium, 293
Symphyodon, 433+; americanus, 433+; machrisianus, 433+, 434*
Syrrhopodon **cyrtacanthos** sp. nov., 95*; laevis, 88+; prolifer, 88; stonae, 88

TALBOT, S. S., S. LOOMAN TALBOT & J. W. THOMSON. Lichens of Tuxedni Wilderness Area, Alaska, 20
TALBOT, S. L. *See* TALBOT, S. S., S. L. TALBOT & J. W. THOMSON, 20
Tanzania, Adelothecium bogotense new from, 287
Tasmania, Plagiothecium novae-seelandiae new from, 221; Siphulella coralloidea new from, 186
Taxiphyllum laevifolium, 435; machrisianum, 433
teloschistin, 153+
Tephromela, 430; aglaea, 27; armeniaca, 403
Tetraphidaceae, 443+
Tetraphis, 174+; pellucida, 174+, 324
Tetrodontium repandum, 443+
Texosporium, 329+; sancti-jacobi, 329+, 331*
Thamnolia subuliformis, 28, 403, 458; vermicularis, 28, 403, 458
thamnolic acid, 458
Thelidium minutulum, 403
Thelocarpon, 227; coccophorum, *228*
Thelomma mammosum, 441; ocellatum, 33+
Thelotrema, 233
THIERS, B. M. & K. S. GILES EMORY. The history of bryology in California, 68
thiophanic acid, 53
THOMSON, J. W. [Review:] H. Schöller. Flechtenverbreitung und Klima. Vegetationsökologische Untersuchungen zur Rolle der Flechten in naturnahen Traubeneichenwäldern des Taunus, 488
THOMSON, J. W. & W. A. WEBER. Lichens collected on the Arctic Excursion of the 9th International Botanical Congress (Montreal) in 1959, 392
THOMSON, J. W. *See* TALBOT, S. S., S. L. TALBOT & J. W. THOMSON, 20
THOR, G. *See* GOWARD, T. & G. THOR, 33
THRALL, P. H. *See* MISHLER, B. D., P. H. THRALL, J. S. HOPPLE, JR., E. DE LUNA & R. VILGALYS, 172
Thuidium, 175+; delicatulum, 174; furfurosum var. sparsum, 9
Thysananthus, 42+; sect. Convolutae, 42; sect. Spathulistipae, 42; sect. Vittatae, 43; aculeatus, 44; amazonicus, 43+; australis, 43+; comosus, 42+; convolutus, 44; dissopterus, 42; evansii, 43+; flavescens, 43; fruticosus, 43+; minor, 44+; mollis, 43+; planus, 43; pterobryoides, 43+; retusus, 43+; spathulistipus, 43+
Timmia austriaca, 31
TIXIER, P. Deux espèces nouvelles Malgaches du genre Calymperopsis (Musci), 284
Tomentypnum nitens, 409+
Toninia aromatica, 403; lobulata, 403; sedifolia, 403; squalida, 403
TONSBERG, T. Rinodina sheardii, a new lichen species from northwest Europe and northwest North America, 216
TONSBERG, T. & I. M. BRODO. Enterographa zonata and Opegrapha gyrocarpa new to North America, 225
TORRENTE, P. *See* EGEA, J. M. & P. TORRENTE, 161
Tortula, 263; andicola, 263; bartramii, 74; bolanderi, 74; pagorum, 9+; papillosa, 9, 263; princeps, 9+; revolvens, 280
Trachyphyllum 206; dusenii, 293
Trachypus bicolor, 289
Trapelia placodioides, 36
Trapeliopsis flexuosa, 28; granulosa, 28
Tremolecia atrata, 403
Trentepohlia, 228
2,4,5-trichlorolichexanthone, 53
2,5,7-trichlorolichexanthone, 52+
2,5,7-trichloro-3-*O*-methylnorlichexanthone, 53
4,5,7-trichloro-6-*O*-methylnorlichexanthone, 52+
Trichostomopsis australasiae, 75; fayae, *75*
Trichostomum brachydontium, 9+
1,3,6-tri-*O*-methylarthothelin, 53+, 54*
Triquetrella papillata, 9+
Tristichium mirabile, 293
Tritomaria quinquedentata, 146, 268
Tuckermannopsis chlorophylla, 28; ciliaris, 403; **commixta** comb. nov., 403; **hepatizon** comb. nov., 403; oakesiana, 139, 167; pinastri, 28, 404; sepincola, 28, 404
Tylimanthus ruwenzorensis, 289

Umbilicaria, 316, 380; cinereorufescens, 28, 404; cylindrica, 28, 404; deusta, 404; esculenta, 316+; hyperborea, 18, 28, 404; mammulata, 167, 316+, 404; muhlenbergii, 404; phaea, 18, 28; polyphylla, 404; proboscidea, 404; torrefacta, 404; vellea, 28
Umbilicariaceae, 316, 380
Underwood, L. M., 72+
uronic acid, 181
Usnea florida, 28; herrei, 139, cf. strigosa, 167
usnic acid, 377+

VAN ROOY, J. & A. E. VAN WYK. A conspectus of the subfamily Macromitrioideae (Bryopsida: Orthotrichaceae) in southern Africa, 205
VAN WYK, A. E. *See* VAN ROOY, J. & A. E. VAN WYK, 205

Verrucaria aethiobola, 28; arctica, 404; devergens, 404; deversa, 404; margacea, 404; nigrescens, 404
Vesicularia, 334
Vestergrenopsis isidiata, 404
VILGALYS, R. *See* MISHLER, B. D., P. H. THRALL, J. S. HOPPLE, JR., E. DE LUNA & R. VILGALYS, 172
vinetorin, 53
Vittia pachyloma, 293

WARNER, B. G. *See* ARAVENA, R. & B. G. WARNER, 445
WATKINSON, S. C. & F. WATT. The cellular localisation of elements in leaf cells of Sphagnum with the scanning proton microprobe, 181
Watson, S., 70
WATT, F. *See* WATKINSON, S. C. & F. WATT, 181
WEBER, W. A. *See* THOMSON, J. W. & W. A. WEBER, 392
Weissia brachycarpa, 11; controversa, 9+; willisiana, 11
Whipple, A. W., 76
Whipple Expedition, 69
WHITTEMORE, A. T. [Review:] R. R. Ireland and L. M. Ley. Atlas of Ontario Mosses, 372
WHITTEMORE, A. T. *See* L. R. STARK & A. T. WHITTEMORE, 65
Wilkes Expedition, 69
WRIGHT, D. Thamnolia (Ascomycotina: Lichenes Imperfecti): First find for California and correction of published mapping of the genus, 458

xanthones, 52+
Xanthoparmelia, 15+, 378+; alectoronica, 378; coloradoensis, 330; conspersa, 139, 378; distincta, *384*; plittii, 18, 139, 167
Xanthoria, 233; **alaskana** sp. nov., 28*; candelaria, 29, 404; elegans, 29, 404; polycarpa, 29; sorediata, 29, 404
Xylographa abietina, 29, 404; elegans, 157; vitiligo, 404

YANO, O. & H. CRUM. Sphagnum tabuleirense, a new species from Brazil, 343

zeorin, 217
Zygodon, 263; pichinchensis, 263+
Zygodontoideae, 205